



ICD-9-CM Code	ICD-10-CM Code(s)
<b>V72.31</b>	Routine gynecological exam
ICD-10-CM Codes	<b>Z01.411</b> Encounter for gynecological examination (general) (routine) with abnormal findings * Use additional code to identify abnormal findings <b>Z01.419</b> Encounter for gynecological examination (general) (routine) without abnormal findings Instructional note under codes should be as follows: **Use additional code: for screening for human papillomavirus, if applicable ( <b>Z11.51</b> ) for screening vaginal pap smear, if applicable ( <b>Z12.72</b> ) to identify acquired absence of uterus, if applicable ( <b>Z90.71</b> -)
<b>701.0</b>	Circumscribed scleroderma
ICD-10-CM Codes	<b>L94.0</b> Localized scleroderma [morphea] <b>L90.0</b> Lichen sclerosis et atrophicus <b>L94.1</b> Linear scleroderma <b>L94.3</b> Sclerodactyly
<b>624.9</b>	Noninflammatory disorders of the vulva and perineum, unspecified
ICD-10-CM Codes	<b>N90.9</b> Noninflammatory disorder of vulva and perineum, unspecified *There are more specific code choice selections available in ICD-10-CM. These include: <b>N90.0</b> Mild vulvar dysplasia <b>N90.1</b> Moderate vulvar dysplasia <b>N90.3</b> Dysplasia of vulva, unspecified <b>N90.4</b> Leukoplakia of vulva <b>N90.5</b> Atrophy of vulva <b>N90.6</b> Hypertrophy of vulva <b>N90.7</b> Vulvar cyst <b>N90.810</b> Female genital mutilation status, unspecified <b>N90.811</b> Female genital mutilation status, Type I status <b>N90.812</b> Female genital mutilation status, Type II status <b>N90.813</b> Female genital mutilation status, Type III status <b>N90.818</b> Other female genital mutilation status <b>N90.89</b> Other specified noninflammatory disorders of vulva and perineum
<b>627.1</b>	Postmenopausal bleeding
	<b>N95.0</b> Postmenopausal bleeding
<b>625.9</b>	Female genital symptoms NOS
ICD-10-CM Codes	<b>N94.89</b> Other specified conditions associated with female genital organs and menstrual cycle <b>N94.9</b> Unspecified condition associated with female genital organs and menstrual cycle <b>R10.2</b> Pelvic and perineal pain <b>R19.8</b> Other specified symptoms and signs involving the digestive system and abdomen <b>R39.9</b> Unspecified symptoms and signs involving the genitourinary system
<b>644.03</b>	Threatened preterm labor, antepartum (without delivery)
ICD-10-CM Codes	<b>O47.00</b> False labor before 37 weeks of completed gestation, unspecified trimester <b>O47.02</b> False labor before 37 completed weeks of gestation, second trimester <b>O47.03</b> False labor before 37 completed weeks of gestation, third trimester

ICD-9-CM Code	ICD-10-CM Code(s)
ICD-10-CM Codes (cont.)	<b>O47.1</b> False labor at or after 37 completed weeks of gestation <b>O47.9</b> False labor, unspecified ** An additional code from category <b>Z3A</b> is needed to specify the weeks of gestation of the pregnancy.
<b>618.01</b>	Cystocele, midline
	<b>N81.11</b> Cystocele, midline
<b>626.2</b>	Excessive menstruation
ICD-10-CM Codes	<b>N92.0</b> Excessive and frequent menstruation with regular cycle <b>N92.1</b> Excessive and frequent menstruation with irregular cycle <b>N92.2</b> Excessive menstruation at puberty <b>N92.4</b> Excessive bleeding in the premenopausal period
<b>648.33</b>	Drug dependence, antepartum
ICD-10-CM Codes	<b>O99.320</b> Drug use complicating pregnancy, unspecified trimester <b>O99.321</b> Drug use complicating pregnancy, first trimester <b>O99.322</b> Drug use complicating pregnancy, second trimester <b>O99.323</b> Drug use complicating pregnancy, third trimester ** The codes from this subcategory require an additional code from <b>F11-F16</b> and <b>F18-F19</b> to identify manifestations of the drug use. ** An additional code from category <b>Z3A</b> is needed to specify the weeks of gestation of the pregnancy.
<b>627.3</b>	Postmenopausal atrophic vaginitis
	<b>N95.2</b> Postmenopausal atrophic vaginitis
<b>646.83</b>	Other specified complications of pregnancy, antepartum
ICD-10-CM Codes	<b>O26.891</b> Other specified pregnancy related conditions, first trimester <b>O26.892</b> Other specified pregnancy related conditions, second trimester <b>O26.893</b> Other specified pregnancy related conditions, third trimester <b>O26.899</b> Other specified pregnancy related conditions, unspecified trimester * There are other code selections available for codes previously coded to <b>646.83</b> : <b>O26.10</b> Low weight gain in pregnancy, unspecified trimester <b>O26.11</b> Low weight gain in pregnancy, first trimester <b>O26.12</b> Low weight gain in pregnancy, second trimester <b>O26.13</b> Low weight gain in pregnancy, third trimester <b>O26.40</b> Herpes gestationis, unspecified trimester <b>O26.41</b> Herpes gestationis, first trimester <b>O26.42</b> Herpes gestationis, second trimester <b>O26.43</b> Herpes gestationis, third trimester <b>O26.811</b> Pregnancy related exhaustion and fatigue, first trimester <b>O26.812</b> Pregnancy related exhaustion and fatigue, second trimester <b>O26.813</b> Pregnancy related exhaustion and fatigue, third trimester <b>O26.819</b> Pregnancy related exhaustion and fatigue, unspecified trimester **An additional code from category <b>Z3A</b> is needed to specify the weeks of gestation of the pregnancy.
<b>632</b>	Missed abortion
ICD-10-CM Codes	<b>O02.1</b> Missed abortion ** Use additional code from category <b>O08</b> to identify any associated complication ** An additional code from category <b>Z3A</b> is needed to specify the weeks of gestation of the pregnancy.



ICD-9-CM Code	ICD-10-CM Code(s)
<b>616.10</b>	Vaginitis and vulvovaginitis, unspecified
ICD-10-CM Codes	<b>N76.0</b> Acute vaginitis **Use additional code ( <b>B95-B97</b> ), to identify infectious agent There are other codes from this subcategory to describe vaginitis: <b>N76.1</b> Subacute and chronic vaginitis <b>N76.2</b> Acute vulvitis <b>N76.3</b> Subacute and chronic vulvitis <b>N76.89</b> Other specified inflammation of vagina and vulva
<b>618.00</b>	Unspecified prolapse of vaginal walls
ICD-10-CM Codes	<b>N81.9</b> Female genital prolapse,unspecified *There are more specific code choice selections available in ICD-10-CM. These include: <b>N81.0</b> Urethrocele <b>N81.10</b> Cystocele, unspecified <b>N81.11</b> Cystocele, midline <b>N81.12</b> Cystocele, lateral <b>N81.2</b> Incomplete uterovaginal prolapse <b>N81.3</b> Complete uterovaginal prolapsed <b>N81.4</b> Uterovaginal prolapsed, unspecified <b>N81.5</b> Vaginal enterocoele <b>N81.6</b> Rectocele <b>N81.81</b> Perineocele <b>N81.82</b> Incompetence or weakening of pubocervical tissue <b>N81.83</b> Incompetence or weakening of rectovaginal tissue <b>N81.84</b> Pelvic muscle wasting <b>N81.85</b> Cervical stump prolapsed <b>N81.89</b> Other female genital prolapsed
<b>620.2</b>	Ovarian cyst NOS/NEC
ICD-10-CM Codes	<b>N83.20</b> Unspecified ovarian cysts <b>N83.29</b> Other ovarian cysts * There are more specific code choice selections available in ICD-10-CM. These include: <b>N83.1</b> Corpus luteum cyst <b>N83.0</b> Follicular cyst of ovary <b>D27.9</b> Benign neoplasm of unspecified ovary <b>Q50.1</b> Developmental ovarian cyst <b>O34.80</b> Maternal care for other abnormalities of pelvic organs, unspecified trimester <b>O34.81</b> Maternal care for other abnormalities of pelvic organs, first trimester <b>O34.82</b> Maternal care for other abnormalities of pelvic organs, second trimester <b>O34.83</b> Maternal care for other abnormalities of pelvic organs, third trimester <b>O65.5</b> Obstructed labor due to abnormality of maternal pelvic organs ** An additional code from category <b>Z3A</b> is needed to specify the weeks of gestation of the pregnancy.
<b>623.9</b>	Noninflammatory disorders of vagina, NOS
ICD-10-CM Codes	<b>N89.8</b> Other specified noninflammatory disorders of vagina * There are more specific code choice selections available in ICD-10-CM. These include: <b>N89.0</b> Mild vaginal dysplasia <b>N89.1</b> Moderate vaginal dysplasia <b>N89.3</b> Dysplasia of vagina, unspecified

ICD-9-CM Code	ICD-10-CM Code(s)
ICD-10-CM Codes (cont.)	<b>N89.4</b> Leukoplakia of vagina <b>N89.5</b> Stricture and atresia of vagina <b>N89.6</b> Tight hymenal ring <b>N89.7</b> Hematocolpos <b>N89.9</b> Noninflammatory disorder of vagina, unspecified
<b>V25.2</b>	Desire for sterilization <b>Z30.2</b> Encounter for sterilization
<b>618.1</b>	Uterine Prolapse w/o mention of vaginal wall prolapse
ICD-10-CM Codes	<b>N81.2</b> Incomplete uterovaginal prolapse <b>N81.3</b> Complete uterovaginal prolapse <b>N81.4</b> Uterovaginal prolapse, unspecified
<b>788.20</b>	Urinary retention, NOS
ICD-10-CM Codes	<b>R33.9</b> Retention of urine, unspecified * There are more specific code choice selections available in ICD-10-CM. These include: <b>R33.8</b> Other retention of urine <b>R33.0</b> Drug induced retention of urine ** Use additional code for adverse effect, if applicable, to identify drug ( <b>T36-T50</b> with fifth or sixth character 5) <b>F45.8</b> Other somatoform disorders
<b>788.31</b>	Urge incontinence
ICD-10-CM Codes	<b>N39.41</b> Urge incontinence ** Code also any associated overactive bladder ( <b>N32.81</b> )
<b>788.33</b>	Mixed incontinence
ICD-10-CM Codes	<b>N39.46</b> Mixed incontinence ** Code also any associated overactive bladder symptoms ( <b>N32.81</b> )
<b>789.03</b>	Abdominal pain RLQ <b>R10.31</b> Right lower quadrant pain
<b>625.6</b>	Female stress incontinence
ICD-10-CM Codes	<b>N39.3</b> Stress incontinence (female) (male) ** Code also any associated overactive bladder symptoms ( <b>N32.81</b> )
<b>V22.1</b>	Supervision other normal pregnancy
ICD-10-CM Codes	<b>Z34.80</b> Encounter for supervision of other normal pregnancy, unspecified trimester <b>Z34.81</b> Encounter for supervision of other normal pregnancy, first trimester <b>Z34.82</b> Encounter for supervision of other normal pregnancy, second trimester <b>Z34.83</b> Encounter for supervision of other normal pregnancy, third trimester
<b>611.72</b>	Lump or mass in breast <b>N63</b> Unspecified lump in breast
<b>599.0</b>	Urinary tract infection, NOS
ICD-10-CM Codes	<b>N39.0</b> Urinary tract infection, site not specified ** Use additional code ( <b>B95-B97</b> ), to identify infectious agent *There are more specific code choice selections available in ICD-10-CM. These include: <b>N30.00</b> Acute cystitis without hematuria <b>N30.01</b> Acute cystitis with hematuria



ICD-9-CM Code	ICD-10-CM Code(s)
<b>599.0</b>	Continued
ICD-10-CM Codes (cont.)	<b>N30.10</b> Interstitial cystitis (chronic) without hematuria <b>N30.11</b> Interstitial cystitis (chronic) with hematuria <b>N30.20</b> Other chronic cystitis without hematuria <b>N30.21</b> Other chronic cystitis with hematuria <b>N30.30</b> Trigonitis without hematuria <b>N30.31</b> Trigonitis with hematuria <b>N30.40</b> Irradiation cystitis without hematuria <b>N30.41</b> Irradiation cystitis with hematuria <b>N30.80</b> Other cystitis without hematuria <b>N30.81</b> Other cystitis with hematuria <b>N30.90</b> Cystitis, unspecified without hematuria <b>N30.91</b> Cystitis, unspecified with hematuria <b>N15.9</b> Renal tubule-interstitial disease, unspecified <b>N34.1</b> Nonspecific urethritis <b>N34.2</b> Other urethritis
<b>627.9</b>	Menopausal disorder, NOS
ICD-10-CM Codes	<b>N95.9</b> Unspecified menopausal and perimenopausal disorder *There are more specific code choice selections available in ICD-10-CM. These include: <b>N95.0</b> Postmenopausal bleeding <b>N95.1</b> Menopausal and female climacteric states <b>N95.2</b> Postmenopausal atrophic vaginitis <b>N95.8</b> Other specified menopausal and perimenopausal disorders
<b>788.34</b>	Incontinence without sensory awareness
ICD-10-CM Codes	<b>N39.42</b> Incontinence without sensory awareness ** Code also any associated overactive bladder ( <b>N32.81</b> )
<b>642.43</b>	Mild or unspecified pre-eclampsia, NOS
ICD-10-CM Codes	<b>O14.00</b> Mild to moderate pre-eclampsia, unspecified trimester <b>O14.02</b> Mild to moderate pre-eclampsia, second trimester <b>O14.03</b> Mild to moderate pre-eclampsia, third trimester <b>O14.90</b> Unspecified pre-eclampsia, unspecified trimester <b>O14.92</b> Unspecified pre-eclampsia, second trimester <b>O14.93</b> Unspecified pre-eclampsia, third trimester **An additional code from category <b>Z3A</b> is needed to specify the weeks of gestation of the pregnancy.
<b>795.01</b>	Papanicolaou smear of cervix with atypical squamous cells of undetermined significance ASC-US
ICD-10-CM Codes	<b>R87.610</b> Atypical squamous cells of undetermined significance on cytologic smear of cervix (ASC-US) *There are more specific code choice selections available in ICD-10-CM. These include: <b>R87.611</b> Atypical squamous cells cannot exclude high grade squamous intraepithelial lesion on cytologic smear of cervix (ASC-H) <b>R87.612</b> Low grade squamous intraepithelial lesion on cytologic smear of cervix (LSIL) <b>R87.613</b> High grade squamous intraepithelial lesion on cytologic smear of cervix (HSIL) <b>R87.614</b> Cytologic evidence of malignancy on smear of cervix <b>R87.615</b> Unsatisfactory cytologic smear of cervix

ICD-9-CM Code	ICD-10-CM Code(s)
<b>795.01</b>	Continued
ICD-10-CM Codes (cont.)	<b>R87.616</b> Satisfactory cervical smear but lacking transformation zone <b>R87.618</b> Other abnormal cytological findings on specimens from cervix uteri <b>R87.619</b> Unspecified abnormal cytological findings in specimens from cervix uteri
<b>626.4</b>	Irregular menstrual cycle
ICD-10-CM Codes	<b>N92.5</b> Other specified irregular menstruation <b>N92.6</b> Irregular menstruation, unspecified * There are more specific code choice selections available in ICD-10-CM. These include: <b>N92.0</b> Excessive and frequent menstruation with regular cycle <b>N92.1</b> Excessive and frequent menstruation with irregular cycle <b>N92.2</b> Excessive menstruation at puberty <b>N92.4</b> Excessive bleeding in the premenopausal period
<b>651.03</b>	Twin pregnancy antepartum
ICD-10-CM Codes	<b>O30.001</b> Twin pregnancy, unspecified number of placenta and unspecified number of amniotic sacs, first trimester <b>O30.002</b> Twin pregnancy, unspecified number of placenta and unspecified number of amniotic sacs, second trimester <b>O30.003</b> Twin pregnancy, unspecified number of placenta and unspecified number of amniotic sacs, third trimester <b>O30.009</b> Twin pregnancy, unspecified number of placenta and unspecified number of amniotic sacs, unspecified trimester <b>O30.011</b> Twin pregnancy, monochorionic/monoamniotic, first trimester <b>O30.012</b> Twin pregnancy, monochorionic/monoamniotic, second trimester <b>O30.013</b> Twin pregnancy, monochorionic/monoamniotic, third trimester <b>O30.019</b> Twin pregnancy, monochorionic/monoamniotic, unspecified trimester <b>O30.031</b> Twin pregnancy, monochorionic/diamniotic, first trimester <b>O30.032</b> Twin pregnancy, monochorionic/diamniotic, second trimester <b>O30.033</b> Twin pregnancy, monochorionic/diamniotic, third trimester <b>O30.039</b> Twin pregnancy, monochorionic/diamniotic, unspecified trimester <b>O30.041</b> Twin pregnancy, dichorionic/diamniotic, first trimester <b>O30.042</b> Twin pregnancy, dichorionic/diamniotic, second trimester <b>O30.043</b> Twin pregnancy, dichorionic/diamniotic, third trimester <b>O30.049</b> Twin pregnancy, dichorionic/diamniotic, unspecified trimester <b>O30.091</b> Twin pregnancy, unable to determine number of placenta and number of amniotic sacs, first trimester <b>O30.092</b> Twin pregnancy, unable to determine number of placenta and number of amniotic sacs, second trimester <b>O30.093</b> Twin pregnancy, unable to determine number of placenta and number of amniotic sacs, third trimester <b>O30.099</b> Twin pregnancy, unable to determine number of placenta and number of amniotic sacs, unspecified trimester **An additional code from category <b>Z3A</b> is needed to specify the weeks of gestation of the pregnancy.



ICD-9-CM Code	ICD-10-CM Code(s)	
<b>795.03</b>	Papanicolaou smear of cervix with LGSIL	
ICD-10-CM Codes	<b>R87.612</b> Low grade squamous intraepithelial lesion on cytologic smear of cervix (LGSIL)	
<b>623.5</b>	Leukorrhea, not specified as infective	<b>N89.8</b> Other specified noninflammatory disorders of vagina
<b>V25.11</b>	Insertion of IUD	<b>Z30.430</b> Encounter for insertion of intrauterine contraceptive device
<b>658.23</b>	Delayed delivery after spontaneous or unspecified rupture of membranes, antepartum condition or complication	
ICD-10-CM Codes	<b>O42.111</b> Preterm premature rupture of membranes, onset of labor more than 24 hours following rupture, first trimester <b>O42.112</b> Preterm premature rupture of membranes, onset of labor more than 24 hours following rupture, second trimester <b>O42.113</b> Preterm premature rupture of membranes, onset of labor more than 24 hours following rupture, third trimester <b>O42.119</b> Preterm premature rupture of membranes, onset of labor more than 24 hours following rupture, unspecified trimester ** An additional code from category <b>Z3A</b> is needed to specify the weeks of gestation of the pregnancy.	
<b>789.00</b>	Abdominal pain site unspecified	
ICD-10-CM Codes	<b>R10.9</b> Unspecified abdominal pain * There are more specific code choice selections available in ICD-10-CM. These include: <b>R10.0</b> Acute abdomen <b>R10.10</b> Upper abdominal pain, unspecified <b>R10.11</b> Right upper quadrant pain <b>R10.12</b> Left upper quadrant pain <b>R10.13</b> Epigastric pain <b>R10.2</b> Pelvic and perineal pain <b>R10.30</b> Lower abdominal pain, unspecified <b>R10.31</b> Right lower quadrant pain <b>R10.32</b> Left lower quadrant pain <b>R10.33</b> Periumbilical pain <b>R10.84</b> Generalized abdominal pain	
<b>V28.3</b>	Encounter for routine screening for malformation using ultrasonics (routine fetal ultrasound, NOS)	<b>Z36</b> Encounter for antenatal screening of mother
<b>V76.2</b>	Screening malignant neoplasm cervix	<b>Z12.4</b> Encounter for screening for malignant neoplasm of cervix
<b>655.13</b>	Maternal care for fetal chromosomal abnormality	
ICD-10-CM Codes	*A 7th character extender must be assigned to code under category <b>O35</b> . <b>O35.1XX0</b> Maternal care for (suspected) chromosomal abnormality in fetus, not applicable or unspecified <b>O35.1XX1</b> Maternal care for (suspected) chromosomal abnormality in fetus, fetus 1 <b>O35.1XX2</b> Maternal care for (suspected) chromosomal abnormality in fetus, fetus 2	

ICD-9-CM Code	ICD-10-CM Code(s)	
ICD-10-CM Codes (cont.)	<b>O35.1XX3</b> Maternal care for (suspected) chromosomal abnormality in fetus, fetus 3 <b>O35.1XX4</b> Maternal care for (suspected) chromosomal abnormality in fetus, fetus 4 <b>O35.1XX5</b> Maternal care for (suspected) chromosomal abnormality in fetus, fetus 5 <b>O35.1XX9</b> Maternal care for (suspected) chromosomal abnormality in fetus, other fetus One of the following 7th characters is to be assigned to each code under category <b>O35</b> . 7th character 0 is for single gestations and multiple gestations where the fetus is unspecified. 7th character 1 through 9 are for cases of multiple gestations to identify the fetus for which the code applies. The appropriate code from category <b>O30</b> , multiple gestation, must also be assigned when assigning a code from category <b>O35</b> that has a 7th character of 1 through 9. 7th Character extenders for subcategory <b>O35</b> - <b>0</b> Not Applicable or unspecified <b>1</b> Fetus 1 <b>2</b> Fetus 2 <b>3</b> Fetus 3 <b>4</b> Fetus 4 <b>5</b> Fetus 5 <b>9</b> Other fetus **An additional code from category <b>Z3A</b> is needed to specify the weeks of gestation of the pregnancy.	
<b>789.04</b>	Abdominal pain LLQ	<b>R10.32</b> Left lower quadrant pain
<b>789.07</b>	Abdominal pain generalized	<b>R10.84</b> Generalized abdominal pain
<b>595.1</b>	Interstitial cystitis chronic	
ICD-10-CM Codes	<b>N30.10</b> Interstitial cystitis (chronic) without hematuria <b>N30.11</b> Interstitial cystitis (chronic) with hematuria	
<b>595.9</b>	Cystitis NOS	
ICD-10-CM Codes	<b>N30.90</b> Cystitis, unspecified without hematuria <b>N30.91</b> Cystitis, unspecified with hematuria **Use additional code to identify infectious agent ( <b>B95-B97</b> ) * There are more specific code choice selections available in ICD-10-CM. These include: <b>N30.00</b> Acute cystitis without hematuria <b>N30.01</b> Acute cystitis with hematuria <b>N30.10</b> Interstitial cystitis (chronic) without hematuria <b>N30.11</b> Interstitial cystitis (chronic) with hematuria <b>N30.20</b> Other chronic cystitis without hematuria <b>N30.21</b> Other chronic cystitis with hematuria <b>N30.30</b> Trigonitis without hematuria <b>N30.31</b> Trigonitis with hematuria <b>N30.40</b> Irradiation cystitis without hematuria <b>N30.41</b> Irradiation cystitis with hematuria <b>N30.80</b> Other cystitis without hematuria <b>N30.81</b> Other cystitis with hematuria	
<b>788.43</b>	Nocturia	<b>R35.1</b> Nocturia
<b>112.1</b>	Candidial vaginitis	<b>B37.3</b> Candidiasis of vulva and vagina
<b>616.2</b>	Bartholin's gland cyst	<b>N75.0</b> Cyst of Bartholin's gland



ICD-9-CM Code	ICD-10-CM Code(s)
<b>616.4</b>	Abcess of vulva NEC <b>N76.4</b> Abscess of vulva
<b>V25.09</b>	Contraceptive management NEC
ICD-10-CM Codes	<b>Z30.09</b> Encounter for other general counseling and advice on contraception <b>Z30.8</b> Encounter for other contraceptive management
<b>659.73</b>	Abnormality in fetal heart rate or rhythm, antepartum
ICD-10-CM Codes	<b>O76</b> Abnormality in fetal heart rate and rhythm complicating labor and delivery ** An additional code from category <b>Z3A</b> is needed to specify the weeks of gestation of the pregnancy.
Many codes from chapter 15 ( <b>O00-O99A</b> ) require the use of an additional code from category <b>Z3A</b> . Weeks of gestation, to identify the specific week of the pregnancy. These codes are for use, only on the maternal record, to indicate the weeks of gestation of the pregnancy.	
ICD-10-CM Codes	Code first complications of pregnancy, childbirth and the puerperium ( <b>O00-O9A</b> ). <b>Z3A.00</b> Weeks of gestation of pregnancy not specified <b>Z3A.01</b> Less than 8 weeks gestation of pregnancy <b>Z3A.08</b> 8 weeks gestation of pregnancy <b>Z3A.09</b> 9 weeks gestation of pregnancy <b>Z3A.10</b> 10 weeks gestation of pregnancy <b>Z3A.11</b> 11 weeks gestation of pregnancy <b>Z3A.12</b> 12 weeks gestation of pregnancy

ICD-9-CM Code	ICD-10-CM Code(s)
ICD-10-CM Codes	<b>Z3A.13</b> 13 weeks gestation of pregnancy <b>Z3A.14</b> 14 weeks gestation of pregnancy <b>Z3A.15</b> 15 weeks gestation of pregnancy <b>Z3A.16</b> 16 weeks gestation of pregnancy <b>Z3A.17</b> 17 weeks gestation of pregnancy <b>Z3A.18</b> 18 weeks gestation of pregnancy <b>Z3A.19</b> 19 weeks gestation of pregnancy <b>Z3A.20</b> 20 weeks gestation of pregnancy <b>Z3A.21</b> 21 weeks gestation of pregnancy <b>Z3A.22</b> 22 weeks gestation of pregnancy <b>Z3A.23</b> 23 weeks gestation of pregnancy <b>Z3A.24</b> 24 weeks gestation of pregnancy <b>Z3A.25</b> 25 weeks gestation of pregnancy <b>Z3A.26</b> 26 weeks gestation of pregnancy <b>Z3A.27</b> 27 weeks gestation of pregnancy <b>Z3A.28</b> 28 weeks gestation of pregnancy <b>Z3A.29</b> 29 weeks gestation of pregnancy <b>Z3A.30</b> 30 weeks gestation of pregnancy <b>Z3A.31</b> 31 weeks gestation of pregnancy <b>Z3A.32</b> 32 weeks gestation of pregnancy <b>Z3A.33</b> 33 weeks gestation of pregnancy <b>Z3A.34</b> 34 weeks gestation of pregnancy <b>Z3A.35</b> 35 weeks gestation of pregnancy <b>Z3A.36</b> 36 weeks gestation of pregnancy <b>Z3A.37</b> 37 weeks gestation of pregnancy <b>Z3A.38</b> 38 weeks gestation of pregnancy <b>Z3A.39</b> 39 weeks gestation of pregnancy <b>Z3A.40</b> 40 weeks gestation of pregnancy <b>Z3A.41</b> 41 weeks gestation of pregnancy <b>Z3A.42</b> 42 weeks gestation of pregnancy <b>Z3A.49</b> Greater than 42 weeks gestation of pregnancy

*Fast Forward*

*to ICD-10*



ICD-10

*Fast Forward*

*to ICD-10*



ICD-10